

[54] ENZYME IMMUNOASSAYS USING
IMMOBILIZED REAGENTS IN A FLOWING
STREAM[75] Inventors: Don N. Gray; Melvin H. Keyes, both
of Sylvania, Ohio[73] Assignee: Technicon Instruments Corporation,
Tarrytown, N.Y.

[21] Appl. No.: 954,269

[22] Filed: Oct. 24, 1978

[51] Int. Cl.³ C12M 1/34; G01N 33/54;
G01N 21/00[52] U.S. Cl. 435/7; 435/291;
23/230 B; 424/12; 422/81[58] Field of Search 435/4, 7, 288, 291,
435/14; 204/195 B; 23/230 B; 422/68, 63, 67,
71, 81, 82, 101; 424/12

[56] References Cited

U.S. PATENT DOCUMENTS

3,902,970	9/1975	Levin	435/14
3,925,018	12/1975	Saunders	435/7
3,975,237	8/1976	Rubenstein	435/7
3,993,451	11/1976	Verbeck	435/14
4,009,005	2/1977	Johnson	422/71
4,022,577	5/1977	Brooker et al.	23/230 B
4,125,492	11/1978	Cuatrecasas et al.	210/31 C
4,141,687	2/1979	Forrest et al.	435/7
4,200,625	4/1980	Reese	23/230 B

OTHER PUBLICATIONS

Ishikawa, "Enzyme Immunoassay of Insulin by Fluorimetry of the Insulin-Glucoamylase Complex," *J. Biochem.*, vol. 73, No. 6, (1973), pp. 1319-1321.Ismail et al., "The South Mend System," A Simple, Fully-Automated, Continuous-Flow System for Immunoassays [Appendix: Application to Serum Thyroxine Radioimmunoassay], *Clin. Chem.*, vol. 24, No. 4, (1978), pp. 571-579.

Brooker et al., "Gammaflow: A Completely Automated

Radioimmunoassay System," *Science*, vol. 194, (1976), pp. 270-276.Primary Examiner—Thomas G. Wiseman
Attorney, Agent, or Firm—Morgan, Finnegan, Pine,
Foley & Lee

[57] ABSTRACT

This invention relates to a method and apparatus for the rapid quantitative determination of an antigen contained in aqueous biochemical samples, for example, blood. The method comprises a series of sequentially arranged stages through which a buffered stream flows. The sample, containing an unknown concentration of the antigen, is injected into the buffered flowing stream and contacts various reagents in the sequential series of stages. An initial solubilization stage comprises an immobilized antibody on a substrate which antibody is specific to the antigen in the sample. The immobilized antibody has been reacted previously to saturation with an enzyme-antigen complex, the antigen of the complex being the same as the antigen in the sample, and the complex is reversably bound to the immobilized antibody. The antigen in the sample and buffered stream flow through the solubilization stage and a competitive equilibrium reaction takes place between the complex bound to the antibody and the antigen in the sample, resulting in the release of a quantity of the enzyme-antigen complex into the flowing stream. The complex and the stream pass into a conversion stage wherein the complex reacts with an immobilized substrate, such as starch, to produce a measurable product which is a quantitative index of the complex released in the solubilization stage. The measurable product and the stream flow to a detection stage wherein the product is reacted, quantitatively measured and related to the concentration of the antigen in the original sample.

24 Claims, 3 Drawing Figures

